

On the universal theory of torsion and lacunary hyperbolic groups

D. Osin *

Abstract

We show that the universal theory of torsion groups is strongly contained in the universal theory of finite groups. This answers a question of Dyson. We also prove that the universal theory of some natural classes of torsion groups is undecidable. Finally we observe that the universal theory of the class of hyperbolic groups is undecidable and use this observation to construct a lacunary hyperbolic group with undecidable universal theory. Surprisingly, torsion groups play an important role in the proof of the latter results.

Keywords: Universal theory, torsion group, lacunary hyperbolic group.

2000 Mathematical Subject Classification: 20F50, 03C60, 20F65, 20F67.

1 Introduction

Recall that a *universal sentence* in a first order language \mathcal{L} is any sentence of the form

$$\forall x_1 \dots \forall x_k \Phi,$$

where Φ is a quantifier free formula. The *universal theory* of a class of groups \mathcal{C} is defined to be the set of all universal sentences in the first order group theoretic language that hold in all groups from \mathcal{C} . As usual, we denote the universal theory of \mathcal{C} by $\text{Th}_\forall(\mathcal{C})$.

Clearly if \mathcal{C} contains another class, say \mathcal{D} , then $\text{Th}_\forall(\mathcal{C}) \subseteq \text{Th}_\forall(\mathcal{D})$. In particular,

$$\text{Th}_\forall(\mathcal{T}) \subseteq \text{Th}_\forall(\mathcal{F}), \tag{1}$$

where \mathcal{T} and \mathcal{F} are the classes of torsion groups and finite groups, respectively. In [4], Dyson asked whether the inclusion in (1) is actually an equality. The positive answer would imply some surprising results, e.g., the absence of finitely presented infinite groups of finite exponent, as observed by Fine, Gaglione, and Spellman [5]. Our first goal is to answer this question.

Theorem 1.1. *We have $\text{Th}_\forall(\mathcal{T}) \subsetneq \text{Th}_\forall(\mathcal{F})$.*

*This work has been supported by the NSF grant DMS-0605093.

Further we show that the universal theory of some natural classes of torsion groups is undecidable. Recall that a theory T in a language \mathcal{L} is *decidable*, if there is an algorithm which, given a sentence Φ in \mathcal{L} , decides whether $\Phi \in T$. In [19], Slobodskoi proved that the universal theory of all finite groups is undecidable. In fact, he also proved that the universal theory of all torsion groups is undecidable although it is not stated explicitly in [19]. We recover the later result and obtain some new ones.

Theorem 1.2. *The universal theory of the following classes of groups is undecidable.*

- a) (Slobodskoi) *The class of torsion groups.*
- b) *The class of groups of any fixed sufficiently large odd non-simple exponent.*
- c) *The class of p -groups for every fixed prime $p \neq 2$.*

Finally we observe that torsion groups can be used to study universal theory of groups of completely different nature, namely hyperbolic and lacunary hyperbolic groups. Recall that a finitely generated group G is *lacunary hyperbolic* if at least one asymptotic cone of G is a real tree [13]. If, in addition, G is finitely presented, then it is hyperbolic. Thus lacunary hyperbolic groups can be thought of as infinitely presented analogues of hyperbolic groups. Alternatively, one may characterize lacunary hyperbolic groups as directed limits of sequences of hyperbolic groups with some additional properties. This characterization is used in [13] to show that lacunary hyperbolic groups resemble hyperbolic ones in many respects.

Recall that every two non-abelian free groups have the same universal theory and this theory is decidable. This solution of the long-standing Tarski Problem was independently obtained by Kharlampovich, Myasnikov [11] and Sela [17]. The question of whether the elementary theory of every hyperbolic group is decidable is still open. A partial result in this direction was obtained by Sela in [18], where he proved that the universal theory of every torsion-free hyperbolic group is decidable. Sela's result was generalized to all hyperbolic groups (possibly with torsion) by Dahmani and Guirardel [3]. The primary goal of our last theorem is to show that there is no hope to generalize it to lacunary hyperbolic groups.

Theorem 1.3. *Let \mathcal{H} denote the class of all hyperbolic groups.*

- 1. $\text{Th}_\forall(\mathcal{H})$ *is undecidable.*
- 2. *There exists a lacunary hyperbolic group G such that $\text{Th}_\forall(G) = \text{Th}_\forall(\mathcal{H})$. In particular, $\text{Th}_\forall(G)$ is undecidable.*

Acknowledgments. I am grateful to Ben Fine, Daniel Groves, Alexander Olshanskii, and Mark Sapir for useful discussions and remarks.

2 Torsion groups

Given a subset $S \subseteq G$ of a group G , we denote by $\langle S \rangle^G$ the smallest normal subgroup of G containing S . Recall that a subgroup $H \leq G$ is a Q -*subgroup* of G (or has the *congruence*

extension property in terminology of [14]) if for every normal subgroup N of H , we have $\langle N \rangle^G \cap H = N$. Equivalently, for every $N \triangleleft H$, the natural homomorphism from H/N to $G/\langle N \rangle^G$ is injective. In what follows we denote by \mathfrak{B}_n the variety of all groups of exponent n , i.e., groups satisfying the identity $X^n = 1$. By $B(m, n)$ we denote the free Burnside group of rank m , i.e., the m -generated free group in \mathfrak{B}_n .

We will use the following theorem of Olshanskii and Sapir proved in [14, Theorem 1.3]. The proof was later simplified by Ivanov [8].

Theorem 2.1 (Olsanskii-Sapir, Ivanov). *For every integer $m > 1$ and every odd sufficiently large n (e.g., $n > 2^{48}$), there exists an embedding of $B(m, n)$ into a finitely presented group G with the following properties.*

- a) $B(m, n)$ is a Q -subgroup of G .
- b) The restriction of the natural homomorphism $\varepsilon: G \rightarrow G/G^n$ to $B(m, n)$ is injective and $\varepsilon(B(m, n))$ is a Q -subgroup of G/G^n .

Theorem 1.1 is a particular case of the following more general result.

Theorem 2.2. *Let \mathcal{C} be a class of groups such that $\mathfrak{B}_n \subseteq \mathcal{C}$ for some odd $n > 2^{48}$. Then $\text{Th}_\forall(\mathcal{C}) \subsetneq \text{Th}_\forall(\mathcal{C} \cap \mathcal{F})$.*

Proof. The inclusion $\text{Th}_\forall(\mathcal{C}) \subseteq \text{Th}_\forall(\mathcal{C} \cap \mathcal{F})$ is obvious, so we only need to show that $\text{Th}_\forall(\mathcal{C}) \neq \text{Th}_\forall(\mathcal{C} \cap \mathcal{F})$. Let

$$G = \langle x_1, \dots, x_s \mid R_1, \dots, R_t \rangle \quad (2)$$

be the finitely presented group from Theorem 2.1 that contains $B(2, n)$. Let w be a nontrivial element of $B(2, n)$ whose image in every finite quotient of $B(2, n)$ is trivial. The existence of such an element follows immediately from the solution of the Bounded Burnside Problem by Novikov, Adian (for odd exponents) and Ivanov (in the general case) [1, 8], and the solution of the Restricted Burnside Problem by Zelmanov [20, 21]. We think of w as an element of G and choose a word W in the alphabet $\{x_1^{\pm 1}, \dots, x_s^{\pm 1}\}$ that represents w .

Clearly W represents 1 in every finite quotient of G . Thus the universal sentence

$$L = \forall x_1 \dots \forall x_s (R_1 = 1 \ \& \ \dots \ \& \ R_t = 1 \Rightarrow W = 1)$$

is true in every finite group. Indeed if elements x_1, \dots, x_s of a finite group K satisfy the relations $R_1 = 1, \dots, R_t = 1$, then the subgroup of K generated by x_1, \dots, x_s is a finite quotient of G .

Thus $L \in \text{Th}_\forall(\mathcal{F}) \subseteq \text{Th}_\forall(\mathcal{C} \cap \mathcal{F})$. On the other hand L is not true in G/G^n . Indeed the images of x_1, \dots, x_s under the natural homomorphism $\varepsilon: G \rightarrow G/G^n$ still satisfy the relations $R_1 = 1, \dots, R_t = 1$. However W represents the element $\varepsilon(w)$ in G/G^n , which is nontrivial since the restriction of ε to $B(2, n)$ is injective by Theorem 2.1. Thus $L \notin \text{Th}_\forall(\mathfrak{B}_n)$. As $\mathfrak{B}_n \subseteq \mathcal{C}$ we have $\text{Th}_\forall(\mathcal{C}) \subseteq \text{Th}_\forall(\mathfrak{B}_n)$, and hence $L \notin \text{Th}_\forall(\mathcal{C})$. \square

To prove our next theorem we need a particular case of a result of Kharlampovich [10].

Theorem 2.3 (Kharlampovich). *Suppose that $n = pq$, where $p \geq 3$ is a prime and q has an odd divisor ≥ 665 . Then for some $m \geq 2$ there exists a normal subgroup $N \leq B(m, n)$ such that N is the normal closure of finitely many elements in $B(m, n)$ and the word problem in $B(m, n)/N$ is undecidable.*

Theorem 1.2 is an immediate corollary of the result below.

Theorem 2.4. *Suppose that \mathcal{C} is a class of groups such that $\mathfrak{B}_n \subseteq \mathcal{C}$ for some odd non-prime $n > 2^{48}$. Then $\text{Th}_\forall(\mathcal{C})$ is undecidable.*

Proof. Clearly every non-prime odd $n > 2^{48}$ satisfies the hypothesis of Theorem 2.3. Let N be a normal subgroup of $B(m, n)$, $m \geq 2$, such that N is the normal closure of finitely many elements in $B(m, n)$ and the word problem in $B(m, n)/N$ is undecidable. Further let G be the finitely presented group from Theorem 2.1 that contains $B(m, n)$.

Let A be the natural image of $B(m, n)$ in $G_1 = G/\langle N \rangle^G$. Since $B(m, n)$ is a Q -subgroup of G , we have $A \cong B(m, n)/N$. Note that G_1 is finitely presented as N is the normal closure of finitely many elements in $B(m, n)$. Let us fix some finite presentation

$$G_1 = \langle x_1, \dots, x_s \mid U_1, \dots, U_p \rangle.$$

Similarly let $H = G/G^n$ and $H_1 = H/\langle \varepsilon(N) \rangle^H$, where $\varepsilon: G \rightarrow H$ is the natural homomorphism. Let also B denote the image of $\varepsilon(B(m, n))$ in H_1 . Since the restriction of ε to $B(m, n)$ is injective and $\varepsilon(B(m, n))$ is a Q -subgroup of H we have $B \cong \varepsilon(B(m, n))/\varepsilon(N) \cong B(m, n)/N$. Moreover, it is easy to see that there exists a homomorphism $\gamma: G_1 \rightarrow H_1$ which induces an isomorphism $A \rightarrow B$.

Let y_1, \dots, y_m be generators of A , Y_1, \dots, Y_m words in the alphabet $\mathcal{X} = \{x_1^{\pm 1}, \dots, x_s^{\pm 1}\}$ that represent these elements in G_1 . Given a word $V = V(y_1, \dots, y_m)$ in the alphabet $\mathcal{Y} = \{y_1^{\pm 1}, \dots, y_m^{\pm 1}\}$, let $\widehat{V}(x_1, \dots, x_s)$ denote the word obtained from V by replacing y_i with Y_i for every $i = 1, \dots, m$. For every word V in \mathcal{Y} , let

$$L(V) = \forall x_1 \dots \forall x_s (U_1 = 1 \ \& \ \dots \ \& \ U_p = 1 \Rightarrow \widehat{V}(x_1, \dots, x_s) = 1).$$

Note that if $L(V) \in \text{Th}_\forall(\mathcal{C})$, then L is true in H_1 as $H_1 \in \mathfrak{B}_n \subseteq \mathcal{C}$. Therefore $\widehat{V}(\gamma(x_1), \dots, \gamma(x_s)) = 1$ in B . Since γ is injective on A , $\widehat{V}(x_1, \dots, x_s) = 1$ in A and hence $V = 1$ in A . On the other hand, if $L(V) \notin \text{Th}_\forall(\mathcal{C})$, then there exists a group $C \in \mathcal{C}$ such that $L(V)$ is false in C . This means that there exists a homomorphism $\alpha: G_1 \rightarrow C$ such that $\widehat{V}(\alpha(x_1), \dots, \alpha(x_s)) \neq 1$ in C . Clearly this implies $V \neq 1$ in A .

Thus if we were able to decide if $L(V) \in \text{Th}_\forall(\mathcal{C})$ for every word V in the alphabet \mathcal{Y} , we would be able to solve the word problem in $A \cong B(m, n)/N$. Since the latter problem is undecidable, $\text{Th}_\forall(\mathcal{C})$ is undecidable as well. \square

3 Hyperbolic groups

The following result is an immediate consequence of methods of [19], although it is not stated explicitly there.

Lemma 3.1 (Slobodskoi). *Let T be any theory such that*

$$\text{Th}_\forall(\mathcal{T}) \subseteq T \subseteq \text{Th}_\forall(\mathcal{F}). \quad (3)$$

Then T is undecidable.

Proof. In [18], Slobodskoi constructed an effective sequence of formulas $\Psi(k)$, $k \in \mathbb{N}$, and two (disjoint) recursively inseparable subsets $X, Y \subseteq \mathbb{N}$ such that $\Psi(k)$ holds for every torsion group whenever $k \in X$ and $\Psi(k)$ does not hold in some finite group whenever $x \in Y$. Recall that two disjoint subsets $X, Y \subseteq \mathbb{N}$ are said to be *recursively inseparable* if there is no recursive subset $I \subseteq \mathbb{N}$ such that $X \subseteq I$ and $I \cap Y = \emptyset$.

Let now T be a decidable theory that satisfies (3). Set $I = \{k \in \mathbb{N} \mid \Psi(k) \in T\}$. Since T is decidable, I is recursive. Clearly (3) implies $X \subseteq I$ and $I \cap Y = \emptyset$, which contradicts recursive inseparability of X and Y . \square

The next lemma is quite elementary. In fact, it follows from Malstsev's Local Theorem (see, e.g., [11, Theorem 27.3.3]) applied to groups considered as algebraic systems with one ternary predicate instead of multiplication. We provide a proof for convenience of the reader.

Lemma 3.2. *Let G be a group and \mathcal{C} is a class of groups. Suppose that at least one of the following conditions holds.*

- (a) *There exists a sequence of normal subgroups $N_1 \supseteq N_2 \supseteq \dots$ of G such that $\bigcap_{i=1}^{\infty} N_i = \{1\}$ and $G/N_i \in \mathcal{C}$ for all $i \in \mathbb{N}$.*
- (b) *There is a group G_0 and a sequence of normal subgroups $N_1 \subseteq N_2 \subseteq \dots$ of G_0 such that $G = G_0 / \bigcup_{i=1}^{\infty} N_i$ and $G_0/N_i \in \mathcal{C}$ for every $i \in \mathbb{N}$.*

Then $\text{Th}_\forall(\mathcal{C}) \subseteq \text{Th}_\forall(G)$.

Proof. First assume that (a) holds. Let \mathcal{U} be a non-principal ultrafilter, $P = \prod_{i=1}^{\infty} (G/N_i) / \mathcal{U}$ the corresponding ultraproduct of (G/N_i) 's. It is straightforward to verify that the map $\varepsilon: G \rightarrow P$ defined by $\varepsilon(g) = (gN_1, gN_2, \dots)$ is a homomorphism and $\text{Ker}(\varepsilon) = \bigcap_{i=1}^{\infty} N_i = \{1\}$. Suppose that a first order formula Φ is true in G/N_i for all $i \in \mathbb{N}$. By the Łoś Theorem Φ is true in P . If, in addition, Φ is universal, then it is obviously true in every subgroup of P . In particular, Φ is true in $\varepsilon(G) \cong G$.

Suppose now that (b) holds. Let \mathcal{U} be a non-principal ultrafilter, $\varepsilon: G_0 \rightarrow \prod_{i=1}^{\infty} (G_0/N_i)/\mathcal{U}$ the map defined by $\varepsilon(g) = (gN_1, gN_2, \dots)$. It is easy to verify that ε is a homomorphism and $\text{Ker}(\varepsilon) = \bigcup_{i=1}^{\infty} N_i$. Hence $\varepsilon(G_0) \cong G$. The rest of the proof is the same as in the first case. \square

Recall that a geodesic metric space M is δ -hyperbolic if for every geodesic triangle Δ in M , every side of Δ belongs to the union of the closed δ -neighborhoods of the other two sides. A group G generated by a set X is δ -hyperbolic relative to X if its Cayley graph with respect to X is a δ -hyperbolic metric space. Further let ε be a homomorphism from the group G to another group. The *injectivity radius* of ε with respect to X , denoted $IR_X(\alpha)$, is defined to be the supremum of all $r > 0$ such that ε is injective on a ball of radius r in G (with respect to the word metric corresponding to X). In particular, $IR_X(\alpha) = \infty$ if ε is injective. We will make use of the following characterization of lacunary hyperbolic groups proved in [13, Theorem 1.1].

Theorem 3.3 (Olshanskii, Osin, Sapir). *A finitely generated group G is lacunary hyperbolic if and only if G is the direct limit of a sequence of finitely generated groups and epimorphisms $G_0 \xrightarrow{\varepsilon_1} G_1 \xrightarrow{\varepsilon_2} \dots$ such that G_i is generated by a finite set S_i , $\varepsilon_i(S_i) = S_{i+1}$, each G_i is δ_i -hyperbolic with respect to S_i , and $\delta_i = o(IR_{S_i}(\varepsilon_i))$ as $i \rightarrow \infty$.*

The construction of the lacunary hyperbolic group in Theorem 1.3 is based on the techniques developed in [16], where the methods suggested by Gromov [6] and elaborated by Olshanskii [12] for hyperbolic groups are generalized to relatively hyperbolic groups. We heavily rely on results from [2, 15, 16] and refer to these papers for more details.

Let K be a group hyperbolic relative to a collection of subgroups $\{H_\lambda\}_{\lambda \in \Lambda}$. A subgroup $L \leq K$ is *suitable* if it is not virtually cyclic, is not conjugate to a subgroup of one of H_λ 's, and does not normalize any nontrivial finite normal subgroup of K . This definition is different from the original one from [16], but is equivalent to the latter by [2, Proposition 3.4]. Parts (a)-(d) of the following theorem are proved in [16, Theorem 2.4]. The last part follows immediately from [16, Lemma 5.1] and the proof of [16, Theorem 2.4].

Theorem 3.4. *Let K be a group hyperbolic relative to a collection of subgroups $\{H_\lambda\}_{\lambda \in \Lambda}$, L a suitable subgroup of K , t_1, \dots, t_m arbitrary elements of K , and $S \subseteq K$ a finite subset. Then there exists an epimorphism $\eta: K \rightarrow \overline{K}$ such that:*

- (a) *The restriction of η to $\bigcup_{\lambda \in \Lambda} H_\lambda$ is injective.*
- (b) *The group \overline{K} is hyperbolic relative to $\{\eta(H_\lambda)\}_{\lambda \in \Lambda}$.*
- (c) *For any $i = 1, \dots, m$, we have $\eta(t_i) \in \eta(H)$.*
- (d) *$\eta(L)$ is a suitable subgroup of \overline{K} .*
- (e) *The restriction of η to S is injective.*

Proposition 3.5. *There exists a lacunary hyperbolic group G such that every hyperbolic group embeds in G .*

Proof. Since every hyperbolic group is finitely presented, we can enumerate all hyperbolic groups: $H_0 = \{1\}, H_1, H_2, \dots$. We set $G_0 = F(x, y)$, where $F(x, y)$ is the free group freely generated by x and y , and construct a sequence of groups and epimorphisms

$$G_0 \xrightarrow{\varepsilon_1} G_1 \xrightarrow{\varepsilon_2} \quad (4)$$

by induction. Suppose that G_n is already constructed. We use the same notation for elements x, y their images in G_n . Assume that the groups H_0, \dots, H_n properly embed in G_n , so we may think of them as subgroups of G_n . Suppose also that G_n is not virtually cyclic, contains no nontrivial finite normal subgroups, and is hyperbolic relative to the collection $\{H_0, \dots, H_n\}$. In particular, G_n is hyperbolic by [15, Corollary 2.41]. These conditions obviously hold for G_0 .

Let δ_n be the hyperbolicity constant of G_n with respect to the generating set $\{x, y\}$, S_n the subset of all elements of G_n of length at most 2^{δ_n} with respect to $\{x, y\}$. Let also $K = G_n * H_{n+1}$. It follows from the definition of relative hyperbolicity via isoperimetric functions (see [15]) that K is hyperbolic relative to $\{H_0, \dots, H_{n+1}\}$. Standard arguments show that $L = G_n$ is a suitable subgroup of K . Let t_1, \dots, t_k be generators of H_{n+1} .

We denote by G_{n+1} the quotient group of K that satisfies the conclusion of Theorem 3.4. By part (a) of Theorem 3.4 we may think of H_0, \dots, H_{n+1} as subgroups of G_{n+1} . Further by part (b) G_{n+1} is hyperbolic relative to $\{H_0, \dots, H_{n+1}\}$. By part (c) the images of t_i 's in G_{n+1} belong to the image of G_n . Thus G_{n+1} is generated by $\{x, y\}$ and there is a natural epimorphism $\varepsilon_{n+1}: G_n \rightarrow G_{n+1}$. Note that by part (e) the natural homomorphism $K \rightarrow G_{n+1}$ is injective on S_n . This implies that

$$IR_{\{x, y\}}(\varepsilon_{n+1}) \geq 2^{\delta_n} \quad (5)$$

Finally by part (d) G_{n+1} is a suitable subgroup of itself. In particular, it is non-elementary and contains no nontrivial finite normal subgroups. This completes the inductive step.

Let now G be the direct limit of (4). That is, let $N_i = \text{Ker}(\varepsilon_n \circ \dots \circ \varepsilon_1)$, $i = 1, 2, \dots$, and $G = G_0 / \bigcup_{i=1}^{\infty} N_i$. Then G is lacunary hyperbolic by (5) and Theorem 3.3. It is clear from our construction that every hyperbolic group embeds in G . \square

Proof of Theorem 1.3. 1) It easily follows from results of [12] that every hyperbolic group is residually torsion. In fact, every hyperbolic group G is even residually torsion of bounded exponent, that is, $\bigcap_{n=1}^{\infty} G^n = \{1\}$, where G^n is the subgroup generated by $\{g^n \mid g \in G\}$. The later result is proved in [9].

Let $N_i = \bigcap_{n=1}^i G^n$. Then the sequence $\{N_i\}$ satisfies the first condition from Lemma 3.2 for $\mathcal{C} = \mathcal{T}$. Hence $\text{Th}_{\forall}(\mathcal{T}) \subseteq \text{Th}_{\forall}(G)$ for every hyperbolic group G . Consequently $\text{Th}_{\forall}(\mathcal{T}) \subseteq \text{Th}_{\forall}(\mathcal{H})$. On the other hand we have $\text{Th}_{\forall}(\mathcal{H}) \subseteq \text{Th}_{\forall}(\mathcal{F})$ as every finite group is hyperbolic. Therefore, $\text{Th}_{\forall}(\mathcal{H})$ is undecidable by Lemma 3.1.

2) Let G be the group from Proposition 3.5. Since every hyperbolic group embeds in G , we have $\text{Th}_\forall(G) \subseteq \text{Th}_\forall(\mathcal{H})$. On the other hand, $\text{Th}_\forall(\mathcal{H}) \subseteq \text{Th}_\forall(G)$ for every lacunary hyperbolic group G by Lemma 3.2 and Theorem 3.3. Thus $\text{Th}_\forall(G) = \text{Th}_\forall(\mathcal{H})$. \square

References

- [1] S.I. Adian, The Burnside problem and identities in groups, *Ergebnisse der Mathematik und ihrer Grenzgebiete* **95**, Springer-Verlag, Berlin-New York.
- [2] G. Arzhantseva, A. Minasyan, D.V. Osin, The SQ-universality and residual properties of relatively hyperbolic groups, *J. Algebra*, **315** (2007), no. 1, 165–177.
- [3] F. Dahmani, V. Guirardel, Foliations for solving equations in groups: free, virtually free, and hyperbolic groups, arXiv:0901.1830.
- [4] V.H. Dyson, A reduction of the open sentence problem for finite groups, *Bull. London Math. Soc.* **13** (1981), no. 4, 331–338.
- [5] B. Fine, A. Gaglione, D. Spellman, Finitely presented infinite torsion groups and a question of V. H. Dyson. In: Combinatorial group theory, discrete groups, and number theory, 119–123, *Contemp. Math.*, 421, Amer. Math. Soc., Providence, RI, 2006
- [6] M. Gromov, Hyperbolic groups. In: *Essays in Group Theory*, MSRI Series, Vol.8, (S.M. Gersten, ed.), Springer, 1987, 75–263.
- [7] S.V. Ivanov, The free Burnside groups of sufficiently large exponents, *Internat. J. Algebra Comput.* **4** (1994), no. 1-2.
- [8] S.V. Ivanov, Embedding free Burnside groups in finitely presented groups, *Geom. Dedicata* **111** (2005), 87–105.
- [9] S.V. Ivanov, A.Yu. Olshanskii, Hyperbolic groups and their quotients of bounded exponents, *Trans. Amer. Math. Soc.* **348** (1996), no. 6, 2091–2138.
- [10] O. Kharlampovich, The word problem for the Burnside varieties, *J. Algebra* **173** (1995), no. 3, 613–621.
- [11] O. Kharlampovich, A. Myasnikov, Elementary theory of free non-abelian groups, *J. Algebra* **302** (2006), no. 2, 451–552.
- [12] A.Yu. Olshanskii, On residualizing homomorphisms and G -subgroups of hyperbolic groups, *Int. J. Alg. Comp.*, **3** (1993), 4, 365–409.
- [13] A. Olshanskii, D. Osin, M. Sapir, Lacunary hyperbolic groups, to appear in *Geom. Top.*; arXiv: math.GR/0701365.
- [14] A. Olshanskii, M. Sapir, Non-amenable finitely presented torsion-by-cyclic groups, *Publ. Math. Inst. Hautes Etudes Sci.* **96** (2002), 43–169.

- [15] D. Osin, Relatively hyperbolic groups: Intrinsic geometry, algebraic properties, and algorithmic problems, *Memoirs Amer. Math. Soc.* **179** (2006), no. 843.
- [16] D. Osin, Small cancellations over relatively hyperbolic groups and embedding theorems, arXiv: math.GR/0411039.
- [17] Z. Sela, Diophantine geometry over groups VI: The elementary theory of a free group, *GAF* **16** (2006), 707-730.
- [18] Z. Sela, Diophantine geometry over groups VII: The elementary theory of a hyperbolic group, *Proceedings of the LMS*, to appear.
- [19] A.M. Slobodskoi, Undecidability of the universal theory of finite groups, *Algebra and Logic* **20** (1981), no. 2, 139–156.
- [20] E.I. Zelmanov, Solution of the restricted Burnside problem for groups of odd exponent, *Math. USSR-Izv.* **36** (1991), no. 1, 41–60.
- [21] E.I. Zelmanov, Solution of the restricted Burnside problem for 2-groups, *Math. USSR-Sb.* **72** (1992), no. 2, 543–565.

Stevenson Center 1326, Department of Mathematics, Vanderbilt University
Nashville, TN 37240, USA

E-mail address: `denis.osin@gmail.com`